

Bipolar Mortality

Date	Author	Place	Age	Sample	Follow-up	OR/SMR/MR	Instrument	DX
2001	Osby <sup>3</sup>	Sweden	15+	15386	10	2.6		ICD
2007	Dutta <sup>1</sup>	London, UK	30+	235	18	1.22	OPCRIT/PSE	DSMIV
2007	Laursen <sup>2</sup>	Denmark	24 - 80+	5558959	30	17.1		ICD-8/10

QUARTILES:

1st	1.9
2nd	2.6
3rd	9.8

Selection of studies described in Eaton, Martins, Nestadt, Bienvenu, Clarke, and Alexandre, *Epidemiologic Reviews*, 2008

### Reference List

- (1) Dutta R, Boydell J, Kennedy N, VAN OJ, Fearon P, Murray RM. Suicide and other causes of mortality in bipolar disorder: a longitudinal study. *Psychol Med.* 2007;37(6):839-847.
- (2) Laursen TM, Munk-Olsen T, Nordentoft M, Mortensen PB. Increased mortality among patients admitted with major psychiatric disorders: a register-based study comparing mortality in unipolar depressive disorder, bipolar affective disorder, schizoaffective disorder, and schizophrenia. *J Clin Psychiatry.* 2007;68(6):899-907.
- (3) Osby U, Brandt L, Correia N, Ekbom A, Sparen P. Excess mortality in bipolar and unipolar disorder in Sweden. *Arch Gen Psychiatry.* 2001;58(9):844-850.