

One year Prevalence of Drug Abuse/Dependence

Date	Author	Place	Sample	RR	Instrument	DX	Rate
1992	Bourdon ³	USA-5 sites	20291	68-79	DIS	DSMIII	2.5
1994	Kessler ⁸	USA-NCS	8098	83	CIDI	DSMIIIR	3.6
1996	Grant ⁴	USA-NLAES	42862	97	AUDADIS	DSMIV	1.5
1998	Bijl ²	Netherlands	7076	70	CIDI	DSMIIIR	1.2
2001	Andrews ¹	Australia	10641	60	CIDI	DSMIV	3
2001	Kringlen ⁹	Norway	2066	78	CIDI	DSMIIIR	0.9
2004	Grant ⁵	USA-NESARC	43093	88	AUDADIS	DSMIV	2
2004	Jacobi ⁶	Germany	4181	81	CIDI	DSMIV	0.7
2005	Kessler ⁷	USA-NCSR	9282	71	CIDI	DSMIV	1.8
2006	Medina-Mora ¹⁰	Mexico	5826	76	CIDI	ICD10	0.4
2007	SAMSHA ¹¹	USA-NSDUH	67802	74	.	DSMIV	2.9

QUARTILES:

1 st	1.1
2 nd	1.8
3 rd	2.7

Selection of studies described in Eaton, Martins, Nestadt, Bienvenu, Clarke, and Alexandre, *Epidemiologic Reviews*, 2008

Reference List

- (1) Andrews G, Henderson S, Hall W. Prevalence, comorbidity, disability and service utilization: overview of the Australian National Mental Health Survey. *British Journal of Psychiatry*. 2001;178:145-153.
- (2) Bijl RV, Ravelli A, van Zessen G. Prevalence of psychiatric disorder in the general population: results of the Netherlands mental health survey and incidence study (NEMESIS). *Social Psychiatry and Psychiatric Epidemiology*. 1998;33:587-595.
- (3) Bourdon KH, Rae DS, Locke BZ, Narrow WE, Regier DA. Estimating the prevalence of mental disorders in U.S. adults from the Epidemiologic Catchment Area Survey
2. *Public Health Rep*. 1992;107(6):663-668.
- (4) Grant BF. DSM-IV, DSM-III-R, and ICD-10 alcohol and drug abuse/harmful use and dependence, United States, 1992: a nosological comparison
1. *Alcohol Clin Exp Res*. 1996;20(8):1481-1488.
- (5) Grant BF, Stinson FS, Dawson DA et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2004;61(8):807-816.
- (6) Jacobi F, Wittchen HU, Holting C et al. Prevalence, co-morbidity and correlates of mental disorders in the general population: results from the German Health Interview and Examination Survey (GHS). *Psychol Med*. 2004;34(4):597-611.
- (7) Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005;62(6):617-27.
- (8) Kessler RC, McGonagle KA, Zhao S et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Archives of General Psychiatry*. 1994;51(1):8-19.

- (9) Kringlen E, Torgersen S, Cramer V. A Norwegian psychiatric epidemiological study
1. *Am J Psychiatry*. 2001;158(7):1091-1098.
- (10) Medina-Mora ME, Borges G, Fleiz C et al. Prevalence and correlates of drug use disorders in Mexico
13. *Rev Panam Salud Publica*. 2006;19(4):265-276.
- (11) Substance Abuse and Mental Health Services Administration. *Results from the 2006 National Survey on Drug Use and Health: National Findings* (Office of Applied Studies, NSDUH Series H-32, DHHS Publication No. SMA 07-4293). 2007.
Ref Type: Report