

One-year Prevalence of Panic Disorder

Date	Author	Place	Panic	Instrument	DX	Sample
1988	Bland <sup>4</sup>	Edmonton, Canada	0.7	CIDI	DSM-III	3,258
1989	Hwu <sup>13</sup>	Taipei, Taiwan	0.2	DIS		5,005
1989	Oakley-Browne <sup>22</sup>	Christchurch, New Zealand	1.4	CIDI	DSM-IV	1,498
1991	Eaton <sup>6</sup>	4 ECA sites	0.9	DIS	DSM-III	14,263
1994	Kessler <sup>15</sup>	US	2.3	CIDI	DSM-IV	8,098
1996	Offord <sup>24</sup>	Ontario, Canada	1.1	CIDI	DSM-III	9,953
1997	Faravelli <sup>7</sup>	Florence, Italy	3.2	CIDI	DSM-III-R	673
1998	Bijl <sup>3</sup>	Netherlands	2.2	CIDI	DSM-III-R	7,076
2001	Andrews <sup>2</sup>	Australia	1.1	CIDI	DSM-IV	10,641
2001	Kringlen <sup>18</sup>	Oslo, Norway	2.6	CIDI	DSM-III-R	2,066
2002	McConnell <sup>20</sup>	Derry, Northern Ireland	2.4	SCAN	DSM-IV	923
2004	Alonso	Belgium, France, Germany, Italy, the Netherlands, and Spain	0.8	CIDI	DSM-IV	21,425
2004	Jacobi <sup>14</sup>	Germany	2.3	CIDI		4,181
2005	Kessler <sup>16</sup>	US	2.7	CIDI	DSM-IV	9,282
2005	Medina-Mora <sup>21</sup>	Mexico	0.6	CIDI	ICD-10	5,826
2005	Pirkola <sup>25</sup>	Finland	1.9	CIDI	ICD-10	6,005
2006	Grant <sup>8</sup>	US	2.1	SADS	DSM-IV	43,093
2006	Kringlen <sup>17</sup>	rural Norway	1.2	CIDI	DSM-III-R	1,714
2006	Shen <sup>27</sup>	metropolitan China	0.2	CIDI	DSM-III-R	5,201
2006	Vicente <sup>28</sup>	Chile	0.9	CIDI	DSM-IV	2,978
2006	Wells <sup>29</sup>	New Zealand	1.7	CIDI	DSM-III-R	12,992
2007	Cho <sup>5</sup>	Korea	0.2	CIDI	DSM-III	6,275
2008	Bromet	Ukraine	1.4	DIS	DSM-IV	1720
2008	Girolamo	Italy	0.7	CIDI	DSM-IV	1779

2008	Gureje <sup>9</sup>	Nigeria	0.1	CIDI	DSM-IV	2,143
2008	Haro <sup>10</sup>	Spain	0.6	CIDI	DSM-IV	2,121
2008	Herman <sup>11</sup>	South Africa	0.8	CIDI	DSM-IV	4,351
2008	Huang <sup>12</sup>	China	0.2	CIDI	DSM-IV	1628
2008	levinson <sup>19</sup>	Israel	0.6	CIDI	DSM-III-R	4859
2008	Oakley-Browne <sup>23</sup>	New Zealand	1.7	CIDI	DSM-III	12,790
2008	Posada-Villa <sup>26</sup>	Colombia	0.7	CIDI	DSM-IV	2,381

---

QUARTILES:

1st	0.6
2nd	0.9
3rd	1.9

---

Selection of studies described in Eaton, Martins, Nestadt, Bienvenu, Clarke, and Alexandre, *Epidemiologic Reviews*, 2008

## Reference List

- (1) Alonso J, Angermeyer MC, Bernert S et al. Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatr Scand Suppl.* 2004;(420):21-27.
- (2) Andrews G, Sanderson K. Prevalence and severity of mental health-related disability and relationship to diagnosis. *Psychiatric services (Washington, D C).* 2001;(178):145-153.
- (3) Bijl RV, Ravelli A, van Zessen G. Prevalence of psychiatric disorder in the general population: results of the Netherlands mental health survey and incidence study (NEMESIS). *Social Psychiatry and Psychiatric Epidemiology.* 1998;33:587-595.
- (4) Bland RC, Newman SC, Orn H. Period prevalence of psychiatric disorders in Edmonton. *Acta Psychiatrica Scandinavica.* 1988;77(Supplement 338):33-42.
- (5) Cho MJ, Kim JK, Jeon HJ et al. Lifetime and 12-month prevalence of DSM-IV psychiatric disorders among Korean adults. *The Journal of Nervous and Mental Disease.* 2007;3(195):203-210.
- (6) Eaton WW, Dryman A, Weissman MM. Panic and Phobia. In: Robins LN, Regier DA, eds. *Psychiatric Disorders in America: The Epidemiologic Catchment Area Study.* New York: The Free Press, 1991: 155-79.
- (7) Faravelli C, Salvatori S, Galassi F, Aiazzi L, Drei C, Cabras P. Epidemiology of somatoform disorders: a community survey in Florence. *Social Psychiatry and Psychiatric Epidemiology.* 1997.
- (8) Grant BF, Hasin DS, Stinson FS et al. The epidemiology of DSM-IV panic disorder and agoraphobia in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry.* 2006;67(3):363-374.
- (9) Gureje O, Adeyemi O, Enyidah N, Ekpo O, Udofia O. Mental Disorders among Adult Nigerians: Risks, Prevalence, and Treatment. In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders.* New York: Cambridge University Press, 2008: 211.

- (10) Haro JM, Alonso J, Pinto-Meza A, Saiz VE, Fernandez A. The Epidemiology of Mental Disorders in the General Population of Spain. In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 406.
- (11) Herman A, Williams D, Stein DJ, Seedat S, Heeringa SG. The South African Stress and Health Study (SASH): A Foundation for Improving Mental Health Care in South Africa . In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 238.
- (12) Huang Y, Liu Z, Zhang M, Shen Y, Tsang A, He Y. Mental Disorders and Service Use in China 447  
*Yueqin Huang, Zhaorui Liu, Mingyuan Zhang, Yucun Shen, Cheuk Him Adley Tsang, Yanling He, and Sing Lee*. In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 447.
- (13) Hwu HG, Yeh EK, Chang LY. Prevalence of psychiatric disorders in Taiwan defined by the Chinese Diagnostic Interview Schedule 1. *Acta Psychiatr Scand*. 1989;79(2):136-147.
- (14) Jacobi F, Wittchen HU, Holting C et al. Prevalence, co-morbidity and correlates of mental disorders in the general population: results from the German Health Interview and Examination Survey (GHS). *Psychol Med*. 2004;34(4):597-611.
- (15) Kessler RC, McGonagle KA, Zhao S et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Archives of General Psychiatry*. 1994;51(1):8-19.
- (16) Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005;62(6):617-27.
- (17) Kringlen E, Torgersen S, Cramer V. Mental illness in a rural area: a Norwegian psychiatric epidemiological study. *Soc Psychiatry Psychiatr Epidemiol*. 2006;41(9):713-719.

- (18) Kringlen E, Torgersen S, Cramer V. A Norwegian psychiatric epidemiological study  
1. *Am J Psychiatry*. 2001;158(7):1091-1098.
- (19) Levinson D, Lerner Y, Zilber N, Levav I, polakiewicz J. The Prevalence of Mental Disorders and Service Use in Israel: Results from the National Health Survey, 2003-2004 . In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 346.
- (20) McConnell P, Bebbington P, McClelland R, Gillespie K, Houghton S. Prevalence of psychiatric disorder and the need for psychiatric care in Northern Ireland. Population study in the District of Derry. *Br J Psychiatry*. 2002;181:214-219.
- (21) Medina-Mora ME, Borges G, Lara C et al. Prevalence, service use, and demographic correlates of 12-month DSM-IV psychiatric disorders in Mexico: results from the Mexican National Comorbidity Survey. *Psychol Med*. 2005;35(12):1773-1783.
- (22) Oakley-Browne MA, Joyce PR, Wells JE, Bushnell JA, Hornblow AR. Christchurch Psychiatric Epidemiology Study, Part II: Six month and other period prevalences of specific psychiatric disorders  
2. *Aust N Z J Psychiatry*. 1989;23(3):327-340.
- (23) Oakley-Browne MA, Wells E, Scott KM. The New Zealand Mental Health Survey 486. In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 486.
- (24) Offord DR, Boyle MH, Campbell D et al. One-year prevalence of psychiatric disorder in Ontarians 15 to 64 years of age  
4. *Can J Psychiatry*. 1996;41(9):559-563.
- (25) Pirkola SP, Isometsä E, Suvisaari J et al. DSM-IV mood-, anxiety- and alcohol use disorders and their comorbidity in the Finnish general population--results from the Health 2000 Study. *Social Psychiatry and Psychiatric Epidemiology*. 2005;1(40):1-10.
- (26) Posada-Villa J, Rodryquez M, Duque P, Garzon A, Breslau J. Mental Disorders in Colombia: Results from the World Mental Health Survey. In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 131.

- (27) Shen YC, Zhang MY, Huang YQ et al. Twelve-month prevalence, severity, and unmet need for treatment of mental disorders in metropolitan China. *Psychol Med.* 2006;36(2):257-267.
- (28) Vicente B, Kohn R, Rioseco P, Saldivia S, Levav I, Torres S. Lifetime and 12-month prevalence of DSM-III-R disorders in the Chile psychiatric prevalence study. *Am J Psychiatry.* 2006;163(8):1362-1370.
- (29) Wells JE, Browne MA, Scott KM, McGee MA, Baxter J, Kokaua J. Prevalence, interference with life and severity of 12 month DSM-IV disorders in Te Rau Hinengaro: the New Zealand Mental Health Survey. *Aust N Z J Psychiatry.* 2006;40(10):845-854.