

One year Prevalence of Depressive Disorder

Date	Author	Place	Sample	RR	Instrument	DX	Rate
1988	Bland ⁶	Edmonton	3258	71.6	DIS	DSMIII	4.6
1989	Hwu ¹⁶	Taipei	5005	95	DIS	DSMIII	0.64
1989	Hwu ¹⁶	Taiwan rural	2995	95	DIS	DSMIII	0.81
1989	Hwu ¹⁶	Taiwan towns	3004	95	DIS	DSMIII	1.1
1989	Oakley-Browne ²⁶	Christchurch	1498	70	DIS	DSMIII	6.7
1990	Faravelli ¹²	Florence	1000	100	SADS	DSMIIR	6.2
1992	Bourdon ⁷	USA-5 sites	20291	68 to 80	DIS	DSMIII	3.5
1994	Kessler ¹⁸	USA	8098	82.4	CIDI	DSMIIR	10.3
1996	Offord ²⁷	Ontario	8116	76.5	CIDI	DSMIIR	4.1
1998	Bijl ⁵	Netherlands	7146	69.7	CIDI	DSMIIR	5.8
1998	Pakriev ²⁹	Udmurtia	855	85.9	CIDI	ICD10/DSMIIR	15.4
1998	Szadoczky ³³	Hungary	2953	85	DIS	DSMIIR	7.1
2000	Lindeman ²²	Finland	5993	86%	UMCIDI-SF	DSMIII	9.3
2001	Andrews ³	Australia	10641	78	CIDI	DSMIV	6.3
2001	Vorcaro ³⁵	Brazil, Bambui	1041	85.3	CIDI	DSMIV	9.1
2002	McConnel ²³	Northern Ireland	1242	87.6	GHQ/SCAN	ICD10	6
2003	Kessler ¹⁹	USA-NCS-R	9090	70.9	WMHCIDI	DSMIV	6.6
2004	ESEMeD ¹¹	Europe-- 6 countries	21425	61.2	WMHCIDI	DSMIV/ICD10	3.9
2004	Wang ³⁶	Canada	17244	na	CIDI-SF	DSMIIR	4.7
2005	Ahola ¹	Finland	3276	88	CIDI	DSMIV	5.2
2005	Beals ⁴	USA, 2 tribes	3084	75.2	CIDI	DSMIIR	3.5
2005	Hasin ¹⁴	USA, NESARC	43093	81	AUDADUS	DSMIV	5.3
2005	Medina-Mora ²⁴	Mexico	5826	76.6	WMHCIDI	DSMIV	3.7
2006	Ohayon ²⁸	South Korea	3719	91.4	SleepEval, phone	DSMIV	3.6
2006	Shen ³¹	China, 2 cities	5201	74.7	WMHCIDI	DSMIV	2
2006	Sloane ³²	Mexico, 4 cities	2509	76.7	CIDI	DSMIV	6.1

2006	Vicente ³⁴	Chile	2978	90.3	CIDI	DSMIIR	5.7
2007	Amoran ²	Oyo, Nigeria	1105	.	GHQ/SCID	DSMIV	5.2
2007	Compton ⁹	USA, NLAES	42862	90	AUDADIS	DSMIV	3.3
2007	Hwang ¹⁵	LA, Chinese	1747	82	CIDI	DSMIV	3.5
2007	Lee ²⁰	Hong Kong	5004	61.2	Telephone	DSMIV	8.4
2007	Williams ³⁷	USA, W	891	69.7	CIDI	DSMIV	6.9
2007	Williams ³⁷	USA, CB	1621	77.7	CIDI	DSMIV	7.2
2007	Williams ³⁷	USA, AA	3578	70.7	CIDI	DSMIV	5.9
2008	Bromet ⁸	Ukraine	1720	78.3	CIDI	DSM-IV	8.4
2008	De Girolamo ¹⁰	Italy	1779	71.3	CIDI	DSM-IV	3
2008	Gureje ¹³	Nigeria	2143	79.3	CIDI	DSM-IV	1.1
2008	Karam ¹⁷	Lebanon	2857	70	CIDI	DSM-IV	4.9
2008	Levinson ²¹	Isreal	4859	72.6	CIDI	DSM-IV	5.9
2008	Oakley-Browne ²⁵	New Zealand	12992	73.3	WMHCIDI	DSMIV	5.7
2008	Posada-Villa ³⁰	Colombia	2381	87.7	CIDI	DSM-IV	5.3
2008	Williams ³⁸	South Africa	4351	85.5	WHOCIDI	DSMIV	4.9

QUARTILES:

1st	3.6
2nd	5.3
3rd	6.5

Selection of studies described in Eaton, Martins, Nestadt, Bienvenu, Clarke, and Alexandre, *Epidemiologic Reviews*, 2008

Reference List

- (1) Ahola K, Honkonen T, Isometsa E et al. The relationship between job-related burnout and depressive disorders--results from the Finnish Health 2000 Study. *J Affect Disord.* 2005;88(1):55-62.
- (2) Amoran O, Lawoyin T, Lasebikan V. Prevalence of depression among adults in Oyo State, Nigeria: a comparative study of rural and urban communities
3. *Aust J Rural Health.* 2007;15(3):211-215.
- (3) Andrews G, Henderson S, Hall W. Prevalence, comorbidity, disability and service utilization: overview of the Australian National Mental Health Survey. *British Journal of Psychiatry.* 2001;178:145-153.
- (4) Beals J, Manson SM, Whitesell NR et al. Prevalence of major depressive episode in two American Indian reservation populations: unexpected findings with a structured interview
4. *Am J Psychiatry.* 2005;162(9):1713-1722.
- (5) Bijl RV, Ravelli A, van Zessen G. Prevalence of psychiatric disorder in the general population: results of the Netherlands mental health survey and incidence study (NEMESIS). *Social Psychiatry and Psychiatric Epidemiology.* 1998;33:587-595.
- (6) Bland RC, Newman SC, Orn H. Period prevalence of psychiatric disorders in Edmonton. *Acta Psychiatrica Scandinavica.* 1988;77(Supplement 338):33-42.
- (7) Bourdon KH, Rae DS, Locke BZ, Narrow WE, Regier DA. Estimating the prevalence of mental disorders in U.S. adults from the Epidemiologic Catchment Area Survey
2. *Public Health Rep.* 1992;107(6):663-668.
- (8) Bromet EJ, Gulzman SF, Tintle NL, Paniotto VI, Webb CP. The State of Mental Health and Alcoholism in Ukraine . In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders.* New York: Cambridge University Press, 2008: 431.
- (9) Compton WM, Thomas YF, Stinson FS, Grant BF. Prevalence, correlates, disability, and comorbidity of DSM-IV drug abuse and dependence in the United States: results from the national epidemiologic survey on alcohol and related conditions
42. *Arch Gen Psychiatry.* 2007;64(5):566-576.

- (10) de Girolamo G, Morosini P, Gigantesco A, Delmonte S, Kessler R.C. The Prevalence of Mental Disorders and Service Use in Italy: Results from the National Health Survey 2001-2003 . In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 364.
- (11) Alonso J, Angermeyer MC, Bernert S et al. Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatr Scand Suppl*. 2004;(420):21-27.
- (12) Faravelli C, Guerrini DB, Aiazzi L, Incerpi G, Pallanti S. Epidemiology of mood disorders: a community survey in Florence 2. *J Affect Disord*. 1990;20(2):135-141.
- (13) Gureje O, Adeyemi O, Enyidah N, Ekpo O, Udofia O. Mental Disorders among Adult Nigerians: Risks, Prevalence, and Treatment. In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 211.
- (14) Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: results from the National Epidemiologic Survey on Alcoholism and Related Conditions 5. *Arch Gen Psychiatry*. 2005;62(10):1097-1106.
- (15) Hwang WC, Myers HF. Major depression in Chinese Americans: the roles of stress, vulnerability, and acculturation. *Soc Psychiatry Psychiatr Epidemiol*. 2007;42(3):189-197.
- (16) Hwu HG, Yeh EK, Chang LY. Prevalence of psychiatric disorders in Taiwan defined by the Chinese Diagnostic Interview Schedule 1. *Acta Psychiatr Scand*. 1989;79(2):136-147.
- (17) Karam EG, Mneimneh ZN, Karam AN, Fayyad JA, Nasser SC, Dimassi I. Mental Disorders and War in Lebanon . In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 265.
- (18) Kessler RC, McGonagle KA, Zhao S et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Archives of General Psychiatry*. 1994;51(1):8-19.

- (19) Kessler RC, Berglund P, Demler O et al. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003;289(23):3095-105.
- (20) Lee S, Tsang A, Zhang MY et al. Lifetime prevalence and inter-cohort variation in DSM-IV disorders in metropolitan China. *Psychol Med*. 2007;37(1):61-71.
- (21) Levinson D, Lerner Y, Zilber N, Levav I, polakiewicz J. The Prevalence of Mental Disorders and Service Use in Israel: Results from the National Health Survey, 2003-2004 . In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 2008: 346.
- (22) Lindeman S, Hamalainen J, Isometsa E et al. The 12-month prevalence and risk factors for major depressive episode in Finland: representative sample of 5993 adults
15. *Acta Psychiatr Scand*. 2000;102(3):178-184.
- (23) McConnell P, Bebbington P, McClelland R, Gillespie K, Houghton S. Prevalence of psychiatric disorder and the need for psychiatric care in Northern Ireland. Population study in the District of Derry. *Br J Psychiatry*. 2002;181:214-219.
- (24) Medina-Mora ME, Borges G, Lara C et al. Prevalence, service use, and demographic correlates of 12-month DSM-IV psychiatric disorders in Mexico: results from the Mexican National Comorbidity Survey. *Psychol Med*. 2005;35(12):1773-1783.
- (25) Oakley-Browne MA, Wells JE, Scott KM. Te Rau Hinengaro: The New Zealand Mental Health Survey. In: Kessler RC, Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York, NY: Cambridge University Press, 2008: 486-508.
- (26) Oakley-Browne MA, Joyce PR, Wells JE, Bushnell JA, Hornblow AR. Christchurch Psychiatric Epidemiology Study, Part II: Six month and other period prevalences of specific psychiatric disorders
2. *Aust N Z J Psychiatry*. 1989;23(3):327-340.
- (27) Offord DR, Boyle MH, Campbell D et al. One-year prevalence of psychiatric disorder in Ontarians 15 to 64 years of age
4. *Can J Psychiatry*. 1996;41(9):559-563.
- (28) Ohayon MM, Hong SC. Prevalence of major depressive disorder in the general population of South Korea. *J Psychiatr Res*. 2006;40(1):30-36.

- (29) Pakriev S, Vasar V, Aluoja A, Saarma M, Shlik J. Prevalence of mood disorders in the rural population of Udmurtia. *Acta Psychiatr Scand.* 1998;97(3):169-174.
- (30) Posada-Villa J, Rodryguez M, Duque P, Garzon A, Breslau J. Mental Disorders in Colombia: Results from the World Mental Health Survey. In: Kessler R.C., Ustun TB, eds. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders.* New York: Cambridge University Press, 2008: 131.
- (31) Shen YC, Zhang MY, Huang YQ et al. Twelve-month prevalence, severity, and unmet need for treatment of mental disorders in metropolitan China. *Psychol Med.* 2006;36(2):257-267.
- (32) Slone LB, Norris FH, Murphy AD et al. Epidemiology of major depression in four cities in Mexico
2. *Depress Anxiety.* 2006;23(3):158-167.
- (33) Szadoczky E, Papp Z, Vitrai J, Rihmer Z, Furedi J. The prevalence of major depressive and bipolar disorders in Hungary. Results from a national epidemiologic survey
1. *J Affect Disord.* 1998;50(2-3):153-162.
- (34) Vicente B, Kohn R, Rioseco P, Saldivia S, Levav I, Torres S. Lifetime and 12-month prevalence of DSM-III-R disorders in the Chile psychiatric prevalence study. *Am J Psychiatry.* 2006;163(8):1362-1370.
- (35) Vorcaro CM, Lima-Costa MF, Barreto SM, Uchoa E. Unexpected high prevalence of 1-month depression in a small Brazilian community: the Bambui Study
1. *Acta Psychiatr Scand.* 2001;104(4):257-263.
- (36) Wang JL. Rural-urban differences in the prevalence of major depression and associated impairment
5. *Soc Psychiatry Psychiatr Epidemiol.* 2004;39(1):19-25.
- (37) Williams DR, Gonzalez HM, Neighbors H et al. Prevalence and distribution of major depressive disorder in African Americans, Caribbean blacks, and non-Hispanic whites: results from the National Survey of American Life
14. *Arch Gen Psychiatry.* 2007;64(3):305-315.
- (38) Williams DR, Herman A, Stein DJ et al. Twelve-month mental disorders in South Africa: prevalence, service use and demographic correlates in the population-based South African Stress and Health Study. *Psychol Med.* 2008;38(2):211-220.